

The intermediate evolution phase in case of truncated selection.

David B. Saakian^{1,2,3}, Christof K. Biebricher⁴, Chin-Kun Hu^{2,5}

¹Yerevan Physics Institute, Alikhanian Brothers St. 2, Yerevan 375036, Armenia,

²Institute of Physics, Academia Sinica, Nankang, Taipei 11529, Taiwan,

³National Center for Theoretical Sciences: Physics Division, National Taiwan University, Taipei 10617, Taiwan,

⁴Max-Planck-Institute for Biophysical Chemistry, D-37070 Göttingen, Germany

⁵Center for Nonlinear and Complex Systems and Department of Physics,
Chung-Yuan Christian University, Chungli 320, Taiwan

(Dated: August 14, 2009)

Using methods of statistical physics, we present rigorous theoretical calculations of Eigen's quasispecies theory with the truncated fitness landscape which dramatically limits the available sequence space of a reproducing quasispecies. Depending on the mutation rates, we observe three phases, a selective one, an intermediate one with some residual order and a completely randomized phase. Our results are applicable for the general case of fitness landscape.

I. INTRODUCTION

Developing realistic evolution models poses an important challenge for evolution research [1, 2]). After the seminal work by Eigen[3] and successful experiments with the self-replication of macromolecules[4, 5], several theoretical studies attempted to explain this molecular evolution phenomenon [6-18]. Realistic fitness landscapes are not smooth, include neutral and lethal types, as observed in recent experimental studies with RNA viruses[20, 21, 22]. In the most of evolution articles symmetric fitness landscapes are considered, where the fitness is a function of the Hamming distance from the wild (reference) sequence. While solving evolution models, the vast majority of results for the mean fitness have been derived using uncontrolled approximations even for the symmetric fitness landscapes, with too simplified sequence space [23] and ignoring back mutations[10]. A simplified geometry with only two (Hamming) classes for sequences with nonzero fitness was used in studies that investigate the role of lethal mutants in evolution[24]. Furthermore, in most evolution models the whole sequence space is assumed to be available for the evolving genome. However, the sequence space a limited population can use is severely restricted to a small part of the sequence space surrounded by an unsurmountable moat of lethal mutations. In this paper, we attempt to rigorously solve this case of a truncated fitness landscape for symmetric fitness landscape.

II. THE SYSTEM

In Eigen's theory[3, 10], an information carrier reproduces with a certain rate r_i , producing offspring of the parental type with the probability Q_{ii} and offspring of another (mutant) type k with the probability Q_{ki} ,

$$dx_i/dt = \{Q_{ii}r_i - \sum_k r_k x_k(t)\}x_i(t) + \sum_{k \neq i} Q_{ik}r_k x_k(t) \quad (1)$$

The whole sequence space contains $M = 4^L$ different sequences, where L is the genome length. The mean fitness of

the system is $R = \sum_k r_k x_k(t)$ where x_i are the relative frequencies of the different genotypes ($\sum_{k=0}^{4^L-1} x_k = 1$). The master type 0 has the maximal fitness r_0 , and its copying fidelity is $Q \equiv Q_{00} = q^L$, where q is the average incorporation fidelity and L the chain length. It is convenient to work with the error rate $u \equiv L(1 - q)$, leading to $Q = e^{-u}$. In this article we consider only the case $L \rightarrow \infty$, u is finite. Eigen maps each genotype precisely into a node on the L -dimensional hypercube and has thus the correct connectivity for each type. The minimal number of steps leading from one position i in sequence space to another one, j , is the Hamming distance d_{ji} .

In the evolution process the information content of the population can be maintained only when the selection force is higher than the dissipating one (mutation). Otherwise, above the *error threshold*, the information gets lost.

It has been shown that the system of nonlinear differential equations in Eq. 1 can be transformed to an infinite system of linear equations, connecting Eigen's model with statistical mechanics[7, 9]. For a single peak fitness landscape ($r_0 = A$ and $r_{i \neq 0} = 1$), the following condition for conserving the master sequence in the population holds [3, 10]:

$$AQ > 1 \quad (2)$$

where $AQ = 1$ is the error threshold.

At the selective phase one has [25]

$$x_0 = \frac{QA - 1}{A - 1} \sim 1, \quad (3)$$

and $x_i \sim 1/L^d$ for $d \ll L$, where d is the Hamming distance from the wild sequence, see Eq. (21) in [15]. We choose the sequences with $1 \leq l \leq L$ from the corresponding l -th Hamming classes. A scaling by Eq.(3) exists also for the rugged (Random Energy Model like) fitness landscapes [11]. Scaling like the one in Eq. (3) has been applied in models of population genetics with few alleles. In realistic fitness landscapes, however, the wild type is present only in a few percents. Assuming neutrality, we can attain such scaling: a substantial fraction of one mutation neighbors of the wild sequence have the same high fitness. Neutrality increases the probability of

such mutants and suppresses x_0 as low as

$$x_0 \sim 1/\sqrt{L}. \quad (4)$$

This result could be derived easily using Eq. (6) in [26], for the case when there is a central neutral sequence and large fracture of neutral sequences among the neighbor sequences of the central sequence.

In non-selective phase one has

$$x_0 \sim 1/M, \quad (5)$$

where M is the total number of sequences, $M = 4^L$.

The error threshold phenomenon closely resembles the ferromagnetic-paramagnetic phase transition[10], where the fitness of the system corresponds to the microscopic energy of the physics system, the mean fitness of the quasiespecies to the free energy, and the mutation rate to temperature. To identify the different phases in statistical physics one uses the free energy and also the order parameters. A phase transition occurs when, during a change of temperature, the analytical expression of the free energy changes. Order parameter changes also: while magnetization is non-zero in the ferromagnetic phase, it is zero at high temperatures and in the absence of a magnetic field. A phase transition in evolution is identified by observing the mean fitness $R = \sum_i x_i r_i$ and choosing proper order parameters, for instance the degree of distribution around the master sequence, the surplus production, $s = \sum_i x_i (1 - 2d_{0i}/L)$, where d_{0i} is the Hamming distance from wild type.

Instead of the 4-letter alphabet of genotypes, we consider only two symbols in a genome, the spins "+" and "-", thus now $M = 2^L$ [8]. Base substitutions correspond to sign changes of the spins. It is particularly easy to analyze landscapes where the fitness values are simple functions of the Hamming distance [8]. The N -dimensional sequence space is then transformed into a quasi-one-dimensional, linear chain of mutant classes l where $p_l = N_l x_l$ comprises all types with the Hamming distance number l from the master and the fitness value J_l . $N_l = \frac{L!}{l!(L-l)!}$ is the number of different genotypes in the class. The parameter l can be identified as a phenotype parameter.

There is a principal difference between quasi-one dimensional model, derived rigorously from the initial sequence space with 2^N sequences and the one-dimensional one considered in [22] and other articles. In contrast to other one-dimensional models used earlier [23] where each class contains only one type, in our case any class l is composed of N_l types and thus retains the connectivity; the Hamming distance between two sequences in the same class can take any value from 0 to $2l$. Moreover, when evolution equations are formulated for class probabilities, the effective mutation rates to the lower class, $\sim (L-l)/L$, and to the higher class, l/L are different and change with l [13],[27]. In contrary, in the one-dimensional model of [22] these mutation rates are l -independent.

In our quasi-one dimensional model J_l can be transformed into the $f(k)$ where $f(k)$ is an appropriate smooth function with the maximum at $k = 1$, and $f(0) = 1$. The "magnetization" parameter k is defined as $k \equiv (1 - 2l/L)$. A correct

version of 1-dimensional evolution model has been suggested first in [27], the discrete time version of parallel model with a linear fitness.

Consider now the solution of the Eigen model with symmetric fitness landscape. The mean fitness R for the fitness function f has been derived as follows: [15]

$$R = \max\{f(k)Q^{(1-\sqrt{1-k^2})}\}_{|-1 \leq k \leq 1}. \quad (6)$$

s can be identified from the mean fitness expression using an equation

$$f(s) = R \quad (7)$$

as has been derived in [13] for the parallel model. Thus in Eq.(6) the maximum is at some k_0 , an order parameter of the system quantifying the bulk spin magnetization, while the surplus s corresponds to the surface magnetization. Eq. (6) is an exact expression (at the infinite genome limit), while in other studies [10, 24] back mutations have been ignored. As shown by Tarazona[9], the Eigen model is not equivalent to the simple ferromagnetic system of spins in the lattice, but only to those spins interacting both inside the bulk of the lattice, and on the surface of the lattice. In this work, different phases will be characterized by R , the mean fitness, by k_0 , the bulk magnetization, by x_0 , the fraction of the wild type of the total population and by s , the surplus. When $k_0 = 0$, resulting in $s = 0$, the population spreads statistically in sequence space, indicating a non-selective phase.

We gave the mean fitness and error threshold (when s in Eq. (7) becomes 0) for the symmetric fitness landscape. The point is that this transition has also information theoretical meaning. Eigen actually found the error threshold from information theoretical consideration of his model. Eigen's idea (information theoretical content of a model) resembles the investigation of information theoretical (optimal coding) aspects of disordered systems, developed in statistical physics two decades later[28, 29]. In the Random Energy Model of spin glass [30] the phase transition point was derived using the information theory analogy [28, 29], and was found to yield results corresponding to those derived by Eigen. The deep information theoretical meaning of error threshold transition in evolution models (equivalent to Shannon inequality for optimal coding) is a solid argument that transition like the one by Eq. (2) exists for any (irregular, with lethal or neutral mutants) fitness landscapes.

III. WAGNER & KRALL THEOREM.

Wagner & Krall [23] considered a population composed of the master and an infinite linear chain of mutants, where each type mutates only to its next neighbor and the fitness r_i decreases monotonically. When there is no low bound of the fitness, an absence of the error threshold transition was derived. Indeed, when in Eq. 6 $f(0) = 0$, there is no error threshold transition. But in more general symmetric fitness landscapes with a finite $f(0)$, this ceases to be valid. The proof is as follows: the maximum of types are located at the Hamming

distance class $L/2$ or, equivalently, at $k = 0$. Consider the logarithm of the right hand side in Eq. 6, and expand near $k = 0$:

$$(1 - \sqrt{1 - k^2}) \ln Q + \ln f(k) \approx -u \frac{k^2}{2} + \ln(f(0)) + ck^\epsilon \quad (8)$$

where c and ϵ are parameters describing the function $f(k)$, and $\ln f(k) - \ln f(0) \sim ck^\epsilon$ at $k \rightarrow 0$. When the fitness decreases slowly and so $\epsilon < 2$, Eq. 8 has a maximum at $k > 0$, fulfilling the condition for selection. When $\epsilon \geq 2$, it can be demonstrated that there is a maximum at $k = 0$ for a sufficiently low reproduction fidelity Q , therefore a sharp error threshold transition results. In the too simplistic model of Wagner & Krall, the right hand side of Eq. 8 lacks the quadratic term, resulting in a monotonic function of k and the absence of phase transition. In the Eigen model the quadratic term holds, breaking the monotonic character of R in Eq. 8 and invoking the error threshold.

IV. TRUNCATED SINGLE PEAK FITNESS LANDSCAPE.

Let us consider a symmetric fitness landscape, where there is non-zero fitness only to some Hamming distance from the reference sequence. Here we define the truncated landscape as a single-peak one where all sequences beyond the Hamming distance $d \equiv L(1 - K)/2$ are lethal:

$$r_l = 1 \quad 1 \leq l \leq d, \quad r_l = 0 \quad l > d \quad (9)$$

Now we have

$$M = \frac{L!}{d!(L - d)!} \quad (10)$$

non-lethal sequences.

To define the mean fitness, we compare the expression of Eq. (6) inside the region $K < k < 1$ and at the border.

The investigation of this model is instructive, see Figs. 1-4. When $QA > 1$ the phase is selective (phase I), x_0 is given by Eq.(3), $k_0 = 1$, $R = QA$, see [31].

When

$$1 > QA > Q^{1-\sqrt{1-K^2}}, \quad (11)$$

a new phase II prevails with

$$k_0 = 1, R = QA \quad (12)$$

In the II phase x_0 is decreasing exponentially with L . The expression of x_0 is calculated in the appendix,

$$x_0 = \exp\left[\int_K^1 dm \frac{1}{2} \ln \frac{\frac{\ln A}{u} + \sqrt{(\frac{\ln A}{u})^2 - 1 + m^2}}{1 + m}\right] \quad (13)$$

At the the transition point (to the II phase) the expression in the exponent is becoming zero, therefore the transition is continuous. When

$$QA < Q^{1-\sqrt{1-K^2}}, \quad (14)$$

the maximum is at the border and we have the non-selective phase III,

$$k_0 = K, R = Q^{1-\sqrt{1-K^2}} \quad (15)$$

The expression for x_0 is defined in the appendix, Eq.(A14). There is some focusing around reference sequence, and x_0 is higher than $1/M$. For $K = 0.9$ we obtained $x_0 \approx 1/M^{2/3}$. The transition between II and III phases is a discontinuous one, x_0 decreases M_1 times, see Eq. (A21)

$$M_1 \sim \exp\left[L \int_K^1 dm \frac{1}{2} \ln \frac{\sqrt{1-K^2} + \sqrt{m^2 - K^2}}{\sqrt{1-K^2} - \sqrt{m^2 - K^2}}\right] \quad (16)$$

Fig.4 illustrates different behavior of x_0 in three phases.

While in the ordinary Eigen model (without truncation of fitness) there is a sharp phase transition with the jump of the x_0 behavior from the Eq.(3) to the $1/M$, in truncated case these sharp transition is moved to the transition point between II and III phases. Now x_0 is continuous at transition point between I and II phases. For the Summers-Litwin case with $K = 1$ we have $M_1 \rightarrow 1$, therefore the transition disappears as has been obtained in [24], and for the $K = 0$ case we get the result of Eigen model $M_1 = 1/2^N$. Our formulas are derived for the case $K \geq 0$.

Is phase II a selective one in the ordinary meaning? Clearly, its mean fitness is higher than in a typical non-selective phase like phase III. This point was clarified by calculating the surplus, replacing the step-like fitness function near the borderline ($k = K$) with a smooth function $f(k)$ which changes its value from 1 to 0 near K . In both phases, II and III, with $QA > Q^{1-\sqrt{1-K^2}}$, the majority of the population is near the borderline, both on the viable and the lethal side. Therefore, while there is a kind of phase transition with some population rearrangement, phase II is identified as an intermediate one, with $x_0 \ll 1$, as in the non-selective phase. Summers and Litwin [24] first realized that $k_0 \rightarrow 1$ in a truncated fitness landscape and tried to analyze the phenomenon. Unfortunately, they used too simplistic a model where all mutants except the nearest neighbors of the master type were lethal. Fig. 2 compares the relative concentration of the master at various superiorities in the single-peak fitness landscape described in [3, 10], with the truncated fitness landscape with $d = 8$, and the case considered by Summers and Litwin. Note the strong dependence of the master concentration on the master superiority in the more realistic landscape, leading to an error threshold, in contrast to the case in [24]. The population profiles of the truncated fitness landscape at different mutation rates are shown in Fig. 3, showing the transition from a master-dominated population via a wide-spread mutant distribution to a non-selective case.

In the Table1 we give the results of numeric for different phases. Mean fitness is well confirmed numerically, while the accuracy of numerics is poor to get correct values of x_0 in the second phase, $2.37 < A < 2.77$.

How the phases can be identified? The fact is that the parameters k_0 and s have different meanings in a statistical physics




FIG. 1: Phase structure for the model with the overlap parameter for truncation point $K = 0.9$, and mutation rate $u = 1$. I: selective phase with $QA > 1$, A is the fitness at the peak, $Q = e^{-u}$ is the errorless copying probability of the genome. II: Intermediate phase with $QA < 1$ and $QA > Q^{1-\sqrt{1-K^2}}$. III: non-selective phase with $QA < Q^{1-\sqrt{1-K^2}}$.




FIG. 2: Semilogarithmic plot of x_0 versus A for a single peak fitness landscape truncated at $d > 8$ with $u = 1$, $L = 3000$. The upper curve corresponds to the Summers-Litwin model, the middle curve to the truncated fitness landscape as in Fig. 3. The square boxes correspond to a single peak fitness model. For $\ln(A) < 1$ Eigen model with single peak fitness has $x_0 = 0$.

approach. This subject has been well analyzed in a series of articles by E. Baake and her co-authors. s and k_0 have been identified with the transverse and longitudinal magnetizations of spins in the corresponding quantum model. We just link s with the mean characteristic of the phenotype, and k_0 with the repertoire of genotypes. The consensus sequence should be determined experimentally not only via distribution x_i , but




FIG. 3: Truncated Eigen model with $d = 8$, $A = 3$, $u = 1$, $L = 3000$. The distribution of probabilities of Hamming classes p_l at different mutation rates (u values). At high accuracy (circles) the population groups around the master type, at intermediate accuracy the population peaks at the truncation border, at low accuracy the master type has practically disappeared and the majority of the progeny is lethal.




FIG. 4: The graphics of $\ln x_0/L$ for the truncated Eigen model at infinite L with $K = 1/2$, $u = 1$. The I phase (curve BC) is at $1 < \ln A/u$, the second (curve DE) at $1 < 0.88 < \ln A/u < 1$ and the third (curve EF) at $0 < \ln A/u < 0.88$. The curve CD corresponds to the jump at the border between II and III phases. In the first phase $\ln x_0/L \sim O(1/L)$.

Phase	III	III	II	II	II	II	I
A	2.2	2.3	2.4	2.5	2.6	2.7	2.8
R	0.859	0.859	0.883	0.919	0.957	0.993	1.030
R th.	0.875	0.875	0.882	0.919	0.956	0.992	1.029
x_0	$4 * 10^{-117}$	$7 * 10^{-54}$	$2 * 10^{-24}$	$4.7 * 10^{-16}$	10^{-9}	$4.4 * 10^{-4}$	0.0167
x_0 th.							0.0164

TABLE I: Eigen model with truncated single peak fitness landscape for $L = 1000$, $u = 1$, $K = 1/2$. The transition between I and second phases is at $A \approx 2.778$, between II and III phases at $A \approx 2.377$.

also via distribution $(x_i)^2 / \sum_{j=0}^M (x_j)^2$. Consider

$$\hat{k}_l = \frac{N_l(x_i)^2}{\sum_j (x_j)^2}, \quad (17)$$

where x_i belongs to the phenotype class l with N_l different genotypes. Having such data, one can simply identify the phase structure.

In the appendix we solve the truncated fitness models for the general monotonic smooth function $f(x)$. The numerics confirms well our analytical results for the new phase.

V. DISCUSSION

We rigorously solved (at the infinite genome length limit) Eigen's model for the truncated selection using the method of [15] as well as methods of statistical mechanics, including the analogy of the error threshold to the ferromagnetic-paramagnetic transition. This analogy is a complicated critical phenomenon, presented by Leuthäusser and Tarazona [7, 9] and well analyzed by E. Baake and co-authors [12, 13]. Instead of using only one order parameter to identify the phase of the model, magnetization, it is necessary to take into account several order parameters describing the order of spins in the bulk lattice and at the surface. Recently the existence of an error threshold has been questioned [24] in the case of truncated selection. In this model the available sequence space has been shrunk to an extremely small size. Fig. 2 illustrates that

the unrealistic assumption in [24] changed the relative concentration of the master type by more than 3 orders of magnitude. Nevertheless, this work was certainly useful for clarifying the concept of quasispecies: the authors first realized the intriguing features of a truncated selection landscape. We found a new evolution (intermediate) phase, when there is no successful selection via phenotype trait (the majority does not share the trait), while there is some grouping of population at genotype level. The intermediate evolution phase differs from the non-selective phase, the frequency of wild type being $\frac{1}{\sqrt{M}}$ or higher in the intermediate phase compared with $\sim \frac{1}{M}$ in the non-selective phase of Eigen model. We proposed a parameter to measure the hidden grouping of population in a genotype level, Eq. (17). Such hidden ordering could be important in case of changing environment: it is possible to force the whole population to extinction changing the fitness of a small fraction (much smaller than $1/L$, but much higher than $1/M$, M is the number of different genotypes) of viruses in the population. We recommend virologists to measure the consensus sequence not only using the probabilities x_i , but also x_i^2 . The evolution picture of the virus population is robust when

two versions of consensus sequence are close to each other. In experiments [21] has been observed an evolution picture, qualitatively similar to the intermediate evolution phase.

How the error threshold transition is connected to the virus extinction in virus experiments, is another story. Several mechanisms are possible: an error catastrophe, as well as a critical mean fitness in order to maintain viral growth [32]. In this work, we observed the new phase with single peak and symmetric landscapes, but this phase exists probably for any (including irregular) fitness landscape with a lethal wall in sequence space.

VI. ACKNOWLEDGMENTS

D.B. Saakian thanks the Volkswagenstiftung grant "Quantum Thermodynamics" and the National Center for Theoretical Sciences in Taiwan and Academia Sinica (Taiwan) under Grant No. AS-95-TP-A07 for the financial support. We thank E. Domingo and M.W. Deem discussions.

-
- [1] E. Domingo, C. K. Biebricher, M. Eigen, and J.J. Holland, *Quasispecies and RNA Virus evolution: Principles and Consequences*. (Landes Bioscience, Austin, TX, 2001).
 - [2] M. Eigen, Proc. Natl. Acad. Sci. USA **99**, 13374 (2002).
 - [3] M. Eigen, Naturwissenschaften **58**, 465 (1971).
 - [4] C. K. Biebricher, Cold Spring Harbor Symp. Quant. Biol. **52**, 299 (1987)
 - [5] D.R. Mills, R. L. Peterson, and S. Spiegelman, Proc. Natl. Acad. Sci. USA **58**, 217 (1967).
 - [6] J. Swetina, and P. Schuster, Biophys. Chem. **16**, 329 (1982).
 - [7] I. Leuthäusser, J. Stat. Phys. **48**, 343 (1987)
 - [8] P. Schuster, and J. Swetina, Bull. Math. Biol. **50**, 635 (1988).
 - [9] P. Tarazona, Phys. Rev. A **45**, 6038 (1992).
 - [10] M. Eigen, J. S. McCaskill, and P. Schuster, Adv. Chem. Phys. **75**, 149 (1989).
 - [11] S. Franz, and L. Peliti, J. Phys. A **30**, 4481 (1997).
 - [12] E. Baake, M. Baake, and H. Wagner, Phys. Rev. Lett. **78**, 559 (1997).
 - [13] E. Baake, and H. Wagner, Genet. Res. **78**, 93 (2001)
 - [14] J. Hermisson, O. Redner, H. Wagner, and E. Baake, Theor. Pop. Biol. **62**, 9 (2002)
 - [15] D. B. Saakian, and C.-K. Hu, Proc. Natl. Acad. Sci. USA **103**, 4935 (2006).
 - [16] D. B. Saakian, E. Munoz, C.-K. Hu, and M. W. Deem, Phys. Rev. E **73**, 041913 (2006).
 - [17] J.-M. Park and M. W. Deem, PRL, **98**, 058101(2007).
 - [18] D.B. Saakian, Journal of statistical physics, **128**,781(2007).
 - [19] D. B. Saakian, O. Rozanova, and Andrei Akmetzhanov, Phys.Rev. E **78**, 041908 (2008).
 - [20] R. Sanjuan, A. Moya, and S. F. Elena, Proc. Natl. Acad. Sci. USA **101**, 8396 (2004).
 - [21] E. Lázaro, C. Escarmis, J. Perez-Mercader, S. C. Manrubia, and E. Domingo, Proc. Natl. Acad. Sci. USA **100**, 10830 (2003).
 - [22] M. Pariera, M., G. Fernandez, B. Clotet, and M. A. Martinez, Mol. Biol. Evol. **24**, 382 (2007).
 - [23] G. P. Wagner, and P. Krall, J. Math. Biol.**32**, 33 (1993).
 - [24] J. Summers, and M. Litwin, J. Virol. **80**, 20 (2006).
 - [25] B. Drossel, Biological evolution and statistical physics Advances in Physics **50**, 209 (2001).
 - [26] E.V. Nimwegen, J.P.Crutchfield, and M. Huynen Proc. Natl. Acad. Sci. USA **96**, 9716 (1999).
 - [27] H. Woodcock and P.G. Higgs, J. Theor. Biol. **179**:61-73, (1996).
 - [28] N. Sourlas, Nature **339**, 693 (1989).
 - [29] D. B. Saakian, JETP Lett.**55**,2(1992).
 - [30] B. Derrida, Phys. Rev. B **24**, 2613 (1981).
 - [31] Kimura and Maruyama. Genetics **54**,1303(1966)
 - [32] J. J. Bull, R. Sanjuan, and C. O. Wilke, J. Virol. **81**, 2930 (2007).

Appendix A. Application of HJE for truncated symmetric landscape.

Let us apply the Hamilton-Jacobi equation (HJE) method [18, 19] to Eigen model with truncated symmetric fitness landscape. Consider a piecewise smooth, monotonic fitness function $f(m)$,

$$\begin{aligned} f(m) &= f_0(m), m > K, \\ f(m) &= 0, m < K \end{aligned} \quad (\text{A.1})$$

Here $f_0(m)$ is a monotonic analytical function.

We denote by k_0 the maximum point in Eq.(6). As s is defined by Eq. (7), for the monotonic fitness function we obtain

$$s \leq k_0, \quad (\text{A.2})$$

Phase	III	III	II	II	II	I	I	I
c	0.5	1	1.2	1.3	1.5	1.9	2.1	3
ln(R)	-0.090	-0.016	0.0163	0.035	0.084	0.212	0.287	0.665
ln(R) th	-0.071	-0.009	0.0160	0.034	0.083	0.213	0.288	0.666
s	0.507	0.507	0.507	0.508	0.510	0.525	0.541	0.663
s th.	0.5	0.5	0.5	0.5	0.500	0.5	0.523	0.666
$-\ln(x_0)$	223	199	185	177	160	131	120	84

TABLE II: Eigen model with truncated fitness, $K = 1/2$, $L = 500$, $u = 1$, $f(x) = \exp(cm^2/2)$. The transition between I and second phases is at $c = 2$, between II and III phases at $c = 1/\sqrt{1-K^2} \approx 1.1547$. In the I phase we have: $s = 1/c$, $\ln(R)/u = c(1-1/c)^2/2$. In the II phase: $s = K$, $\ln(R)/u = c(1-1/c)^2/2$. In the III phase: $s = K$, $\ln(R)/u = cK^2/2 + \sqrt{1-K^2} - 1$.

Phase	III	III	III	II	II	II	I	I
c	0.5	1	2.2	2.4	2.5	2.6	3	4
ln(R)	-0.135	-0.016	-0.022	0.0202	0.0433	0.0676	0.172	0.459
ln(R) th	-0.11	-0.092	-0.042	0.0201	0.0435	0.0678	0.171	0.460
s	0.507	0.507	0.510	0.514	0.518	0.523	0.563	0.698
s th.	0.5	0.5	0.5	0.5	0.5	0.5	0.556	0.701
x_0	230	216	142	128	122	117	98	70

TABLE III: Eigen model with truncated fitness, $K = 1/2$, $L = 500$, $u = 1$, $f(x) = \exp(cm^3)$. The transition point is at $c \approx 2.31$, $m_0 = 0.866$ in the model without truncation. The transition point between II,III is at $c \approx 2.79$.

and there is only one solution of Eq. (7).

Using an ansatz

$$p_l = \exp[LU_0(m, t)], \quad (\text{A.3})$$

$m = 1 - 2l/L$, in [18] has been derived the following equation

$$L \frac{\partial U_0(m, t)}{\partial t} = f_0(m) e^{-u} \times \exp\left\{u \left[\cosh\left(2 \frac{\partial U_0(m, t)}{\partial m}\right) + m \sinh\left(2 \frac{\partial U_0(m, t)}{\partial m}\right) \right]\right\} \quad (\text{A.4})$$

The HJE approach works for any piecewise smooth fitness functions. Eq. (A1) is such a case. To solve our truncated case we should just use different analytical solutions for the regions $-1 < m < K$ and $K < m < 1$.

Assuming an asymptotic $U_0(m, t) = \frac{R}{L}t + U(m)$, we derived [18]

$$R = f_0(m) e^{-u} \times \exp\left\{u \left[\cosh\left(2 \frac{dU(m)}{dm}\right) + m \sinh\left(2 \frac{dU(m)}{dm}\right) \right]\right\}, \quad (\text{A.5})$$

where R is derived by Eq.(6). The surplus s is defined as the value of m where $U(m)$ has a maximum. When s is inside the region $[K, 1]$, $U'(s) = 0$. At extremum points with $U'(m) = 0$ Eq.(A5) gives $f(s) = R$. As for a monotonic fitness function there is a single solution for Eq. (6), $U(x)$ has

a single maximum point in this case, therefore it is a concave function and we take $U(s) = 0$.

We use Eq.(A5) to define p_l with an accuracy $O(1)$ for $\ln p_l$, calculating $U(m) = U(s) + \int_s^m U'(m) dm$ for the corresponding $m = 1 - 2l/L$. Moreover, it is possible to calculate $\ln p_l$ with a higher accuracy $O(1/L)$. In [18] we gave explicit formulas for the case of parallel model. It is possible to construct similar results for the Eigen model as well.

We have two branches of solutions for Eq. (A5):

$$U'(m) = \frac{1}{2} \ln \frac{q \pm \sqrt{q^2 - 1 + m^2}}{1 + m}, \quad q = \frac{1}{u} \ln \frac{R}{f(m)} + 1 \quad (\text{A.6})$$

It is a principal point the choice of different solutions. We choose the proper branch assuming:

- $U(m)$ is continuous function,
- $U'(m)$ is continuous function,
- $U(m)$ is a concave function for the monotonic fitness function $f(m)$.

The transition between two branches ($\pm$ solutions in Eq.(A5)) is only at the point where $q^2 - 1 + m^2 = 0$ or

$$R = f(m) e^{-u+u\sqrt{1-m^2}} \quad (\text{A.7})$$

According to Eq.(6), R is the maximum of the right hand side. Thus we should choose only the branch with " - " sign when k_0 is at the border, $k_0 = K$. When k is inside the interval $[K, 1]$, then we choose the " - " solution for the interval $[k_0, 1]$ and " + " solution in the interval $[K, k_0]$.

For the $V(m) = \ln(\sqrt{N} p_l)/L$ we have another equation [18],

$$R = f_0(m) e^{-u} \exp\left\{u \left[\cosh\left(2 \frac{dV(m)}{dm}\right) \sqrt{1-m^2} \right]\right\} \quad (\text{A.8})$$

The minimum of the right hand side via V' just gives the $f(m) e^{-u+u\sqrt{1-m^2}}$. Thus at the maximum point $m = k_0$ of function $V(m)$ we have $V'(k_0) = 0$. In this article we consider the case when Eq.(6) has a single solution $k = k_0$. Then $V(m)$ is a concave function.

Solution of equations (A5),(A7) are simply related,[18],

$$V(m) = U(m) + \frac{(1+m) \ln \frac{1+m}{2}}{4} + \frac{(1-m) \ln \frac{1-m}{2}}{4} \quad (\text{A.9})$$

Consider now different phases of our model. The selective one with $K < k_0 < 1$, $K < s < 1$; the non-selective one with $k_0 = K$, $s = K$, and intermediate one with $K < k_0 < 1$, $s = K$.

Selective phase.

Now R is given by Eq. (6) with a $K < k_0 < 1$. We used " - " solution of Eq.(A6) for $k_0 < m < 1$ and the " + " solution for $K < m < k_0$. The maximum points of both functions $U(m)$ and $V(m)$ are inside the interval $[K, 1]$. We have $U'(s) = 0$ and $V'(k_0) = 0$. The formulas for the steady

state distributions are the same as in [18]. We have a mean fitness

$$R = f(k_0)e^{-u+u\sqrt{1-k_0^2}} \quad (\text{A.10})$$

For the $p_l, m = L(1 - 2l/L), k_0 < m \leq 1$ we have an expression

$$p_l = \exp[L \int_{k_0}^m dm \frac{1}{2} \ln \frac{q - \sqrt{q^2 - 1 + m^2}}{1 + m} + L \int_s^{k_0} dm \frac{1}{2} \ln \frac{q + \sqrt{q^2 - 1 + m^2}}{1 + m}],$$

$$q = \frac{1}{u} \ln \frac{R}{f(m)} + 1 \quad (\text{A.11})$$

For $m < k_0$ we have

$$p_l = \exp[L \int_s^m dm \frac{1}{2} \ln \frac{q + \sqrt{q^2 - 1 + m^2}}{1 + m}],$$

$$q = \frac{1}{u} \ln \frac{R}{f(m)} + 1 \quad (\text{A.12})$$

Nonselective phase.

Now the maximum of Eq.(6) is at the border $k_0 = K$, and we have

$$R = f(K)e^{-u+u\sqrt{1-K^2}} \quad (\text{A.13})$$

We use " - " solution of Eq.(A6) for the whole interval $K < m < 1$. For the $p_l, m = L(1 - 2l/L)$ we have an expression

$$p_l = \exp[L \int_K^m dm \frac{1}{2} \ln \frac{q - \sqrt{q^2 - 1 + m^2}}{1 + m}],$$

$$q = \frac{1}{u} \ln \frac{R}{f(m)} + 1 \quad (\text{A.14})$$

and the maximum is for p_d with $m \equiv \frac{1-2d}{L} = K$.

For the single peak fitness case ($f(1) = A$ and $f(m) = 1$ for $m < 1$), we have $q = \sqrt{1 - K^2}$ and

$$x_0 \equiv p_0 \sim \exp[\frac{L}{2} \int_K^1 dm \ln \frac{\sqrt{1 - K^2} - \sqrt{m^2 - K^2}}{1 + m}] \quad (\text{A.15})$$

For the $1/M$ we have an expression

$$\frac{1}{M} \sim \exp[\frac{L}{2} \int_K^1 dm \ln \frac{1 - m}{1 + m}] \quad (\text{A.16})$$

Intermediate phase. Now mean fitness is given by Eq.(6) with some $K < k_0 < 1$ and $s = K$. We used " - " solution of Eq.(A6) for $k_0 < m < 1$ and " + " solution for $K < m < k_0$. When $m > k_0$, we have

$$p_l = \exp[L \int_K^{k_0} dm \frac{1}{2} \ln \frac{q + \sqrt{q^2 - 1 + m^2}}{1 + m} + L \int_{k_0}^m dm \frac{1}{2} \ln \frac{q - \sqrt{q^2 - 1 + m^2}}{1 + m}] \quad (\text{A.17})$$

In case of $K < m < k_0$ we have

$$p_l = \exp[L \int_K^m dm \frac{1}{2} \ln \frac{q + \sqrt{q^2 - 1 + m^2}}{1 + m}] \quad (\text{A.18})$$

We took $U(K) = 0$, as the maximum of population is at the border with $m = K$.

For the SP case we have $q = \frac{\ln A}{u}$, and

$$x_0 = \exp[L \int_K^1 dm \frac{1}{2} \ln \frac{\frac{\ln A}{u} + \sqrt{(\frac{\ln A}{u})^2 - 1 + m^2}}{1 + m}] \quad (\text{A.19})$$

Above the transition point $\ln A = \sqrt{1 - K^2}$ Eq.(19) gives

$$x_0 = \exp[L \int_K^1 dm \frac{1}{2} \ln \frac{\sqrt{1 - K^2} + \sqrt{m^2 - K^2}}{1 + m}] \quad (\text{A.20})$$

We see that at the transition point there is a jump, x_0 decreases M_1 times,

$$M_1 \sim \exp[L \int_K^1 dm \frac{1}{2} \ln \frac{\sqrt{1 - K^2} + \sqrt{m^2 - K^2}}{\sqrt{1 - K^2} - \sqrt{m^2 - K^2}}] \quad (\text{A.21})$$

For $K = 0.5$ Eq.(A20) gives $\ln(x_0)/L \approx -0.057$, while $\ln(M)/L \approx -0.56$. Thus above the transition point to the third phase

$$\text{Log}(x_0) \sim \frac{1}{M^{0.1}} \quad (\text{A.22})$$

Consider the case $K = 0.99$. Now we have $\ln(x_0)/L \approx -0.01$ and $\ln(M)/L \approx -0.031$. Thus

$$\ln(x_0) \sim \frac{1}{M^{0.32}} \quad (\text{A.23})$$

Consider evolution model with general fitness function $f(x)$. Assume that without truncation the error threshold transition is a discontinuous one, and there is a jump from non-zero $k_0 > 0$ in selective phase to $k = 0$ solution in Eq.(6) for non-selective phase. Let us introduce the truncation. Choosing $K < k_0$, we have three phases, see Table 3, and x_0 decreases M_2 times at the transition point between II and III phases,

$$M_2 \sim \exp[L \int_{k_0}^1 dm \frac{1}{2} \ln \frac{\sqrt{1 - K^2} + \sqrt{m^2 - K^2}}{\sqrt{1 - K^2} - \sqrt{m^2 - K^2}}] \quad (\text{A.24})$$

If in the original (without truncation) model the error threshold transition is a continuous one with $k_0 = 0$, after truncation we have different expressions for x_0 in the II ($k_0 > K, s = K$) and III ($k_0 = K, s = K$) phases while continuous transitions $I \rightarrow II$ and $II \rightarrow III$, see Table 2. We have a similar behavior for the phase transitions in case of originally (without truncation) discontinuous error threshold transition, if the truncation parameter K is chosen too large, $K > k_0$.

Let us derive an important constraint for the population of the class at the Hamming distance $n = L(1 - k_0)/2$. For the corresponding V we have

$$V(k_0) \equiv \frac{1}{L} \ln \frac{p_n}{\sqrt{N_n}} \quad (\text{A.25})$$

As $p_d = 1$ (the majority of population is at the border with the overlap parameter $K = (1 - 2d)/L$), we have

$$V(K) \equiv \frac{1}{L} \ln \frac{1}{\sqrt{M}}, \quad (\text{A.26})$$

as $M = N_K$. We proved before that $V(m)$ has a single maximum (in our case with a single solution for the maximum point k in Eq. (6)). Thus

$$V(K) > V(k_0) \quad (\text{A.27})$$

which gives

$$p_n > \frac{\sqrt{N_n}}{\sqrt{M}} \quad (\text{A.28})$$

The last inequality supports the choice of order parameter in Eq. (17).

For the single peak case $n = 0$, and we get from Eq.(A28)

$$x_0 > \frac{1}{\sqrt{M}} \quad (\text{A.29})$$

Eqs.(A22,A23) give even higher values for x_0 .